


Family Guide How to use Lyophilised Exosomes



Important

What you receive:

1 Vial Lyophilised Exosomes (10 ml)



+ 1 Vial 30 ml empty and sterile




What you need to buy:



Alcohol pads

Sterile 70% Isopropyl Alcohol, Medical Grade Wipes (individually wrapped); 5 wipes per treatment (one per day);



Sterile Saline Solution

0.9% Sodium Chloride for Inhalation. Either 30ml sterile saline 0.9% or 30ml sterile distilled water;



Syringes and needles

15-20 ml Plastic Syringes with 18G needles, individually sealed and packed. 5 syringes needed per treatment;



Gloves

1 pair of Glove for each Application. 5 pairs in total per treatment;



Vitalia24
INNOVATIVE HEALTHCARE

Important


Preparation




1. Make sure you have a clean work surface away from open windows or doors.




2. Prepare the syringe (10cc - 20cc recommended), alcohol wipes, hospital gloves, sterile fluid, the 10 ml Mesenchymal Stem Cell derived Lyophilised Exosomes and the empty and sterile 30ml vial.



3. Using the syringe take 5ml of sterile solution and inject into the 10 ml vial containing the Lyophilised Exosome powder. Gently turn the vial upside down until the powder is dissolved. Add another 5 ml of sterile solution.



4. Using the syringe extract the 10 ml dissolved exosome solution and inject it into the the sterile 30 ml vial provided.



5. Add 20 ml of sterile solution to the 30 ml vial.

The exosome solution is now ready for administration. Please remember to keep it stored in the fridge.

Important



Vitalia24
INNOVATIVE HEALTHCARE

Application



1. Take the vial with the exosome solution out of the fridge and GENTLY turn it upside down several times in order to have a homogenous solution. Please remember: DO NOT SHAKE!



2. Unpack one syringe with needle.



3. Clean the seal of the vial with the sterile wipe.




4. Insert the needle into the vial and gently pull the syringe to be filled with 5 ml solution.

5. Slowly pull the needle out of the vial and fill the 5 ml solution from the syringe into the chamber of the Nebulizer (if you don't have one yet, you can order it online).

6. Place the Nebulizer mask on the face of the patient and start the inhalation treatment.

7. Repeat this procedure for 5 days (5 ml solution per day).



Additional Info


There is no strict timetable for the inhalation, but do not administer the inhalation for at least 30 minutes AFTER a big meal.



There's no need to hand warm up the bottle.



Try to minimize the time the exosomes are outside of the fridge.



For further questions contact Vitalia24.

